## **Tobacco-Related Cancer Incidence, Excluding Lung Cancer, by Sex, Maine, 2009**

Sex	Count	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine Total	1,058	80.3	75.5 - 85.2	63.5	59.7 - 67.6
Female	369	54.7	49.3 - 60.6	41.1	36.9 - 45.6
Male	689	107.1	99.2 - 115.4	91.9	85.0 - 99.2

## Tobacco-Related Cancer Incidence, Excluding Lung Cancer, by Age, Maine, 2007-2009

Ago	Count	Crude Rate		
Age		Rate	95% CI	
00-14 years	NSD	NSD	NSD	
15-19 years	NSD	NSD	NSD	
20-24 years	NSD	NSD	NSD	
25-34 years*	39	8.7	6.2 - 12.0	
35-39 years	40	15.9	11.4 - 21.7	
40-44 years	89	31.0	24.9 - 38.1	
45-49 years	137	41.7	35.0 - 49.3	
50-54 years	216	66.7	58.1 - 76.2	
55-59 years	324	110.4	98.7 - 123.2	
60-64 years	381	158.8	143.2 - 175.5	
65-69 years	437	249.8	226.9 - 274.4	
75-79 years	426	364.3	330.5 - 400.6	
80-84 years	337	374.4	335.5 - 416.6	
85+ years	331	398.6	356.8 - 443.9	

## **Tobacco-Related Cancer Incidence, Excluding Lung Cancer, by Race, Maine, 2000-2009**

Race	Count	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
White	10,331	81.3	79.8 - 82.9	67.6	66.3 - 69.0
Black	25	18.2	11.8 - 26.9	41.8	26.0 - 62.9
Other**	50	23.8	17.7 - 31.4	45.5	32.0 - 62.3

Data Source: Maine Cancer Registry, November 2011 NPCR data submission

Count: Number of cases diagnosed with malignant cancer.

Tobacco related cancer excluding lung: SEER Cancer site recode: 22020, 20080, 21100, 21010, 21020, 29020, 29010, 27010, 35021 [which include ICD-O-3 codes C320-C329, C100-C109,C150-C159,C160-C169,C250-C259,C530-C539,C670-C679,C649, C659,(C420,

C421, C424, ICD-O-3 histology type:9840, 9861,9866, 9867, 9871-9874, 9895-9897, 9910, 9920]

Crude rates are new cases per 100,000 population

95% CI: 95% confidence interval of the rate.

Rates are calculated using SEER\*Stat Version 7.0.5

NSD: Non-sufficient data. Data is not available due to small or zero counts.

\*The age groups of 25-29 and 30-34 are combined to show sufficient data to calculate a rate for the larger age group.